

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr**SEEBRI® BREEZHALER®**

Glycopyrronium inhalation powder hard capsules

Read this carefully before you start taking **SEEBRI® BREEZHALER®** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **SEEBRI BREEZHALER**.

What is SEEBRI BREEZHALER used for?

- SEEBRI BREEZHALER is used long term in adults to manage airway blockage from chronic obstructive pulmonary disease (COPD; a lung disease that makes it hard to breathe). This can include chronic bronchitis (inflammation of the lungs) and emphysema (damage to parts of the lung known as alveoli).
- SEEBRI BREEZHALER is **not** a rescue medicine and should **not** be used on an as needed basis for treating sudden, severe symptoms of COPD (e.g. sudden breathing problems). Your healthcare professional may give you other medicines to use for sudden breathing problems.

How does SEEBRI BREEZHALER work?

SEEBRI BREEZHALER contains glycopyrronium bromide which belongs to a group of medicines called bronchodilators. In COPD the muscles around the airways tighten, making breathing difficult. SEEBRI BREEZHALER blocks tightening of these muscles in the lungs, making it easier for air to get in and out of the lungs. When you inhale it, it helps you breathe more easily.

What are the ingredients in SEEBRI BREEZHALER?

Medicinal ingredient: glycopyrronium bromide

Non-medicinal ingredients: carrageenan, FDC Yellow 6 (110 Sunset Yellow FCF), hypromellose, lactose monohydrate (which contains milk protein), magnesium stearate, potassium chloride, purified water or

calcium chloride, FDC Yellow 6 (110 Sunset Yellow FCF), hypromellose, lactose monohydrate (which contains milk protein), magnesium stearate

SEEBRI BREEZHALER comes in the following dosage forms:

Inhalation powder hard capsules: 50 mcg of glycopyrronium (as glycopyrronium bromide)

Do not use SEEBRI BREEZHALER if:

- you are allergic to glycopyrronium bromide or any of the other ingredients in SEEBRI BREEZHALER (see **What are the ingredients in SEEBRI BREEZHALER?**)
- you have a severe milk protein allergy as SEEBRI BREEZHALER contains lactose

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take SEEBRI BREEZHALER. Talk about any health conditions or problems you may have, including if you:

- are pregnant or planning to become pregnant
- are breastfeeding or plan to breastfeed
- have or have had heart problems, such as rapid or irregular heartbeat or long QT syndrome
- are taking any medications including eye drops, this includes medications you can buy without prescription
- have difficulty urinating, which can include problems caused by an enlarged prostate
- have eye pain caused by increased pressure in the eyes (narrow-angle glaucoma)
- have kidney problems

Other warnings you should know about:

Eye problems: Avoid getting SEEBRI BREEZHALER powder into your eyes. This may cause eye pain, discomfort, temporary blurring of vision, and/or coloured images in association with red eyes. These may be signs of acute narrow-angle glaucoma (eye pain caused by increased pressure in the eyes). If you develop any of these symptoms, talk to your healthcare professional right away.

Driving and Using Machines: SEEBRI BREEZHALER can cause dizziness or blurred vision. You should not drive or operate machinery if this occurs.

Monitoring: Your healthcare professional might monitor your health throughout your treatment with SEEBRI BREEZHALER. This can include monitoring your kidneys and the development of sudden or worsening COPD symptoms.

Talk to your healthcare professional immediately if:

- your shortness of breath becomes worse
- you don't get the same benefit from your medicine as you did before
- you have breathing difficulties and chest pain
- you have trouble urinating

See the **Serious side effects and what to do about them** table, below, for more information on these and other serious side effects.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with SEEBRI BREEZHALER:

- medicines similar to SEEBRI BREEZHALER (other short- or long-acting muscarinic antagonists) used for your lung disease. They may interact with SEEBRI BREEZHALER and increase the risk of experiencing side effects.

How to take SEEBRI BREEZHALER:

- SEEBRI BREEZHALER has been prescribed for you and should not be given to other people. Take SEEBRI BREEZHALER capsules exactly as your healthcare professional has told you. Do not take more than once a day or exceed the prescribed dose. You should check with your healthcare professional if you are not sure.

- Your healthcare professional may also provide you with an inhaled short-acting bronchodilator for the treatment of COPD symptoms that may occur suddenly.
- The contents of the capsule must be inhaled once daily through the mouthpiece of the SEEBRI BREEZHALER inhalation device only. **Do NOT swallow the capsule.** The SEEBRI BREEZHALER inhaler is especially designed for SEEBRI BREEZHALER capsules and must **not** be used with any other capsules. Likewise, you should not take your SEEBRI BREEZHALER capsules with any inhalation device other than the SEEBRI BREEZHALER inhaler.
- When you start a new pack, use the new SEEBRI BREEZHALER inhaler supplied in the new pack. Dispose of each inhaler after 30 days of use.
- You can inhale SEEBRI BREEZHALER before or after food or drink.
- Store the SEEBRI BREEZHALER capsules in the blister strip until immediately before use.
- Before starting your treatment with SEEBRI BREEZHALER, make sure that you are completely familiar with the use and proper care of the SEEBRI BREEZHALER inhaler. See the **Instructions for Use** for complete information.
- It is important that you continue to take SEEBRI BREEZHALER regularly even if you feel fine and do not have any symptoms.

If you have any questions about SEEBRI BREEZHALER or the SEEBRI BREEZHALER inhaler, talk to your healthcare professional.

Usual dose:

One (1) SEEBRI BREEZHALER capsule is to be inhaled once daily, preferably at the same time each day, only with the SEEBRI BREEZHALER inhaler.


Instructions for Use

This part of the leaflet explains how to use and care for your SEEBRI BREEZHALER inhaler. Please read carefully and follow these instructions.

If you have any questions, ask your healthcare professional.

Your SEEBRI BREEZHALER Inhaler pack contains:

- One SEEBRI BREEZHALER inhaler
- One or more blister cards containing the SEEBRI BREEZHALER capsules to be used in the inhaler



Please read the full “Instructions for Use” before using the SEEBRI BREEZHALER Inhaler.




Insert





Pierce and release






Inhale deeply




Check capsule is empty





Step 1a:
Pull off cap



Step 2a:
Pierce capsule once
Hold the inhaler upright.
Pierce capsule by firmly pressing both side buttons at the same time.
You should hear a noise as the capsule is pierced.
Only pierce the capsule once.



Step 3a:
Breathe out fully
Do not blow into the inhaler.



Check capsule is empty
Open the inhaler to see if any powder is left in the capsule.

If there is powder left in the capsule:

- Close the inhaler.
- Repeat steps 3a to 3c.

Step 1b:
Open inhaler





Step 1c:
Remove capsule
Separate one of the blisters from the blister card.
Peel open the blister and remove the capsule.
Do not push the capsule through the foil.
Do not swallow the capsule.



Step 3b:
Inhale medicine deeply
Hold the inhaler as shown in the picture.
Place the mouthpiece in your mouth and close your lips firmly around it.
Do not press the side buttons.
Breathe in quickly and as deeply as you can.
During inhalation, you will hear a whirring noise.
You may taste the medicine as you inhale.



Step 3c:
Hold breath
Hold your breath for up to 5 seconds.



Remove empty capsule
Put the empty capsule in your household waste.
Close the inhaler and replace the cap.



Step 1d:

Insert capsule

Never place a capsule directly into the mouthpiece.



Step 1e:

Close inhaler

Important Information

- SEEBRI BREEZHALER capsules must always be stored in the blister card and only removed immediately before use.
- Do not push the capsule through the foil to remove it from the blister.
- Do not swallow the capsule.
- Do not use the SEEBRI BREEZHALER capsules with any other inhaler.
- Do not use the SEEBRI BREEZHALER inhaler to take any other capsule medicine.
- Never place the capsule into your mouth or the mouthpiece of the inhaler.
- Do not press the side buttons more than once.
- Do not blow into the mouthpiece.
- Do not press the side buttons while inhaling through the mouthpiece.
- Do not handle capsules with wet hands.
- Never wash your inhaler with water.

Frequently Asked Questions

Why didn't the inhaler make a noise when I inhaled?

The capsule may be stuck in the capsule chamber. If this happens, carefully loosen the capsule by tapping the base of the inhaler. Inhale the medicine again by repeating steps 3a to 3c.

What should I do if there is powder left inside the capsule?

You have not received enough of your medicine. Close the inhaler and repeat steps 3a to 3c.

I coughed after inhaling – does this matter?

This may happen. As long as the capsule is empty you have received enough of your medicine.

I felt small pieces of the capsule on my tongue – does this matter?

This can happen. It is not harmful. The chances of the capsule breaking into small pieces will be increased if the capsule is pierced more than once.

Cleaning the inhaler

Wipe the mouthpiece inside and outside with a clean, dry, lint-free cloth to remove any powder residue. Keep the inhaler dry. Never wash your inhaler with water.

Disposing of the inhaler after use

Each inhaler should be disposed of after all capsules have been used. Ask your pharmacist how to dispose of medicines and inhalers that are no longer required.

Overdose:

If you think you, or a person you are caring for, have inhaled too much SEEBRI BREEZHALER, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Symptoms of an overdose with SEEBRI BREEZHALER include:

- constipation
- difficulty urinating
- increased eye pressure or pain
- eye redness
- vision changes

Missed Dose:

If you miss or forget to take a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the next dose at your regularly scheduled time. Do not take two doses at the same time on the same day.

What are possible side effects from using SEEBRI BREEZHALER?

These are not all the possible side effects you may have when taking SEEBRI BREEZHALER. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- cough
- hoarse voice
- headache
- upset stomach, indigestion
- vomiting
- feeling of pressure or pain in the cheeks and forehead (sinus congestion)
- runny or stuffy nose, sore throat
- dry mouth or throat. Talk to your healthcare professional if the dry mouth persists.
- difficulty sleeping (insomnia)
- fatigue, weakness
- pain in muscles, bones or joints, pain in arms or legs
- tingling or numbness
- dental cavities
- rash, itching
- nose bleeds

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Heart palpitations: unusually fast or irregular heartbeat		✓	
Hyperglycemia (high blood sugar): increased thirst, frequent urination, dry skin, headache, blurred vision and fatigue		✓	
Urinary retention (inability to pass urine or to empty the bladder): difficulty and pain when passing urine, urinating frequently, urination in a weak stream or drips		✓	
Eye problems: new or worsened pressure in your eyes, eye pain or discomfort, blurred vision, seeing halos of bright colours around lights, red eyes			✓
Gastroenteritis (inflammation of the stomach and intestines): abdominal pain, diarrhea, results in nausea, vomiting		✓	
UNKNOWN			

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Paradoxical bronchospasm (sudden narrowing of the airways after taking medicines known as bronchodilators): tightness of the chest associated with coughing, wheezing, or breathlessness immediately after inhalation of SEEBRI BREEZHALER			✓
Allergic reaction: rash, hives, swelling of the face, mouth, lips, throat and tongue, difficulty swallowing or breathing			✓
Cystitis (bladder infection): increased need to urinate, pain in the pelvis or lower back, frequent urination during the night, cloudy urine that may contain blood, burning sensation when passing urine		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Do not use after the expiry date shown on the box.

Store SEEBRI BREEZHALER at room temperature between 15 to 25°C.

Store the capsules in the original package in a dry place in order to protect from heat and moisture. Do not remove capsules from the blister pack until immediately before use.

Do not use this medicine if you notice that the pack is damaged or show signs of tampering.

Each inhaler should be disposed of after 30 days of use.

Keep out of reach and sight of children.

If you want more information about SEEBRI BREEZHALER:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website (www.novartis.ca), Novartis Pharmaceuticals Canada Inc., or by calling 1-800-363-8883.

This leaflet was prepared by Novartis Pharmaceuticals Canada Inc.

Last Revised [October 28, 2022]

Novartis Version: May 26, 2025

SEEBRI and BREEZHALER are registered trademarks.